

PART A2: THE REVOREDO MODEL — TOWARD A THEORETICAL FRAMEWORK FOR THE CONVERSION OF REAL MATTER INTO TACHYONIC MATTER FOR HYPERSPACE TRAVEL

A2.1. Philosophical Foundation

A2.1.1. Why This Model Is Theoretical Physics, Not Science Fiction

The Revoredo Model proceeds from a rigorous epistemological premise: if string theory, the Alcubierre metric, or the ER=EPR conjecture of Maldacena and Susskind are regarded as legitimate theoretical physics despite lacking direct empirical verification, then any hypothesis resting on the same mathematical scaffolding shares an identical epistemological status.

The distinction between science fiction and speculative physics does not lie in current verifiability, but in internal mathematical consistency and the degree of anchorage to formally developed theoretical frameworks. The Revoredo Model satisfies both criteria.

The model does not claim to resolve open problems such as macroscopic quantum decoherence, the extraction of zero-point energy, or the derivation of a unified action from first principles. These are acknowledged as open challenges — precisely the kind of problems that any frontier-physics hypothesis inherits. The Revoredo Model assumes that such barriers could be overcome by a sufficiently advanced civilization, and explores the physical consequences of that assumption without violating known principles. Its epistemic status is therefore not that of a demonstrated theory, but of a mathematically consistent and formally anchored speculative framework, awaiting experimental or observational validation.

That said, the model is not proposed as an abstract exercise in theoretical physics. It is a serious attempt to provide a physically coherent explanation for the anomalous aerial phenomena documented by the U.S. Navy and the Pentagon —

Beyond the Veil: Uncovering the Truth About UFOs

including observations reported in official publications and declassified releases, such as those described by Luis Elizondo and the UAP Task Force. These objects exhibit flight characteristics — extreme acceleration, sudden velocity changes, no visible control surfaces or propulsion signatures — that lie well beyond current human engineering capabilities and push the boundaries of our understanding of physics.

The Revoredo Model offers a candidate mechanism — tachyon-mediated hyperspace transit — that could account for those observables without violating known physical principles. It does not claim that UAPs are necessarily using this mechanism; rather, it claims that if they are physical objects (as the available data suggest), then a framework like this one may be required to explain their behavior. In that sense, the model is a physical hypothesis motivated by empirical observations, not a product of pure speculation.

A2.1.2. The Central Philosophical Argument

Imaginary numbers were considered "unreal" for centuries precisely because they do not appear in the three-dimensional world we perceive. Yet today they are indispensable tools in quantum mechanics for describing subatomic reality.

The argument is straightforward: if the additional dimensions of hyperspace are, by definition, inaccessible to our ordinary three-dimensional perception, then the mathematics governing that domain could well be the mathematics of imaginary numbers. What in our real axis is an "impossible" or "unstable" mass, in the hyperspace axis could be the standard and stable vibration of that environment.

Nature holds its secrets and explains them through mathematics. We must not confuse the local uselessness of a mathematical tool with its ontological unreality.

A2.1.3. The Honest Limits of Current Physics

Contemporary physics is explicitly incomplete on three fronts relevant to this model:

Beyond the Veil: Uncovering the Truth About UFOs

- General Relativity and Quantum Mechanics are mutually incompatible. When one attempts to unify them in extreme regimes (the Big Bang, the interior of black holes), the equations produce infinities, indicating that the theoretical framework is insufficient.
- 95% of the observable universe (dark matter and dark energy) remains without a known causal explanation. It is entirely possible that within that 95% there exist phenomena operating with imaginary-mass mathematics in a stable fashion.
- The light barrier is mathematically strict under the assumption that spacetime is a smooth, continuous fabric—something quantum physics suspects is false at sub-Planckian scales (quantum foam).

A2.2. Mathematical Basis: Tachyons and Special Relativity

A2.2.1. The Relativistic Energy Equation

Relativistic physics defines the total energy (E) of a particle through the fundamental Einstein-Lorentz formula:

$$E = \frac{m_0 c^2}{\sqrt{1 - \frac{v^2}{c^2}}}$$

Where m_0 is the rest mass, v is the particle's velocity, and c is the speed of light in a vacuum.

The Algebraic Extension: Imaginary Masses

If a particle travels faster than light ($v > c$), the term under the square root ($1 - v^2/c^2$) becomes negative. Its square root yields an imaginary number (i). For the energy (E) to be a real, measurable quantity, the rest mass (m_0) must also be a pure imaginary number:

$$m_0 = i \cdot \mu$$

Beyond the Veil: Uncovering the Truth About UFOs

When both imaginary numbers divide in the energy equation, they cancel, and the resulting energy is real. This is the mathematical origin of the tachyon: a particle whose squared mass is negative ($m^2 < 0$).

A2.2.2. The Quantum Field Theory Formulation

In modern physics, tachyons are formalized by introducing a sign change in the squared mass term in wave equations such as the Klein-Gordon equation:

$$(\square + m^2) \varphi = 0 \rightarrow \text{with } m^2 < 0 \text{ for the tachyonic field}$$

This sign change generates an unstable potential with a "Mexican hat" shape. The imaginary mass does not describe a particle that travels backward in time, but a system in an unstable state that will undergo tachyon condensation toward a lower-energy state—a mechanism mathematically identical to that of the Higgs Boson.

A2.3. The Infinite-Energy Barrier

In special relativity, an object with real mass requires ever-increasing energy as it accelerates. To reach $v = c$ would require infinite energy. An object with real mass can never cross the speed of light: it is permanently trapped in the subluminal regime.

The tachyon does not "accelerate" to cross this barrier. Mathematically, it is born directly on the superluminal side ($v > c$) and can never decelerate below the speed of light. The two regimes are separated by an energy singularity, not by a gradual barrier. The reversion follows the same logic: it does not cross c by deceleration either (A2.8.5).

A2.3.1. The Geometric Solution: Hyperspace

To circumvent the infinite-energy problem without violating it, theoretical physics invokes the geometry of hyperspace. In theories such as string theory or Kaluza-Klein spaces, a ship would not accelerate through conventional three-dimensional space, but would jump to a higher dimension. By traveling a

Beyond the Veil: Uncovering the Truth About UFOs

short distance in hyperspace and re-entering normal space, the ship appears to have traveled at superluminal speed, though it locally never exceeded the speed of light.

In the classical model, the ship does not change its material nature: it retains its normal mass. The Revoredo Model goes further.

A2.4. The Tachyonic Conversion Hypothesis

The Revoredo Model proposes unifying the physics of hyperspace with the physics of tachyons under a hypothesis of mass-phase conversion. The process has three structural stages:

- **Mass conversion:** the ship alters its constituent particles, transforming their real mass into imaginary mass ($m^2 < 0$). The particles fundamentally change their ontological state.
- **Barrier jump:** upon becoming imaginary, normal matter instantaneously transforms into tachyons and disappears from conventional three-dimensional space. It does not cross the barrier: it circumvents it by changing its plane of existence.
- **Inverse propulsion:** in the tachyonic world, the energy-velocity relationship is inverted. The less energy a particle has, the faster it travels. Infinite velocity corresponds to zero energy. Spending energy is required to slow down or direct motion.

A2.4.1. The Tachyonic Inversion of Mechanics

In our space: an object at rest has minimum energy ($E = mc^2$). Moving requires gaining kinetic energy.

In tachyonic space: a tachyon traveling at infinite velocity has zero energy. To decelerate to the speed of light requires infinite energy.

The tachyonic ship does not propel itself to advance; it expends energy to orient and decelerate.

A2.4.2. *The Gravitational Bridge of Imaginary Mass*

The tachyonic formulation in the Klein-Gordon equation with $m^2 < 0$ describes the instability of a scalar field, which is mathematically analogous to the Higgs field prior to its spontaneous symmetry breaking. However, establishing that this $m^2 < 0$ produces a gravitational energy density with negative sign in the tensor $T_{\mu\nu}$ requires an intermediate derivation step, which the model closes here.

The energy-momentum tensor of a scalar field φ takes the general form:

$$T_{\mu\nu}^{\varphi} = \partial_{\mu}\varphi \cdot \partial_{\nu}\varphi - g_{\mu\nu} \cdot \left[\frac{1}{2}(\partial\varphi)^2 + V(\varphi) \right]$$

In the standard case of the Higgs field, the system decays spontaneously toward a stable vacuum with positive or zero energy density. The Revored Phase Operator Φ_R introduces the condition that distinguishes the Revored Model from the standard Higgs case:

Revored Inversion Condition (RIC):

$$\Phi_R \text{ active} \leftrightarrow V''(\varphi) < 0 \quad \text{AND} \quad \partial_t(V(\varphi)) > 0$$

The first condition is the negative potential curvature—the standard definition of $m^2 < 0$. The second is the model's innovation: the Revored Engine actively maintains the field within the unstable zone of the potential (from the unstable maximum to the inflection point), through the nonlocal energy compensation of the Revored-EPR Network (A2.7.4), preventing the field from "falling" to the real vacuum. This is an inhibited tachyon condensation. The RIC does not designate a single point but a mode of operation: the engine actively sustains the state of the field against its spontaneous evolution.

Under this condition, the temporal component of the energy-momentum tensor can take a locally negative value—the gravitational requirement for the Revored field equation to operate with local negative energy density. General Relativity does not distinguish the microscopic origin of $T_{\mu\nu}$: it responds

Beyond the Veil: Uncovering the Truth About UFOs

to the full tensor. If the Revoredo Engine sustains a field state with $T_{00} < 0$ locally, geometry responds exactly as the unified field equation requires. The resulting local violation of the Weak Energy Condition is not an absolute prohibition in nature: the Casimir effect and dark energy are physically confirmed examples of its violability in specific regimes.

The formal bridge is closed: the $m^2 < 0$ of Klein-Gordon connects to the negative gravitational $T_{\mu\nu}$ through the mechanism of inhibited tachyon condensation sustained by the Revoredo-EPR Network.

A2.4.3. The Dynamic Equation of Phase Transition: Unification of Sen's Rolling Tachyon with PT-Symmetric Quantum Mechanics

A2.4.3.1. The Ontological Problem of Mass-Phase Conversion

The Revoredo Mechanism requires the transformation of baryonic matter (with real positive mass squared, $m^2 > 0$) into tachyonic matter (with negative mass squared, $m^2 < 0$) without violating unitarity, causality, or local energy conservation. A naïve interpolation in the complex mass plane—as proposed in earlier drafts—constitutes a mathematical description without physical grounding. The correct approach must derive the transition from an existing theoretical framework.

The solution lies in **Sen's rolling tachyon formalism** (Sen 2002a, 2002b), which describes the decay of an unstable D-brane into closed-string tachyonic matter. This formalism is already present in the model's conceptual references (A2.17) and provides the exact dynamic equation for the transition.

A2.4.3.2. Sen's Effective Action as the Foundation

The open-string tachyon field $T(x,t)$, living on the D-brane to which baryonic matter is anchored, is governed by the Dirac–Born–Infeld-type effective action:

$$S_T = - \int d^4x V(T) \sqrt{1 - \eta^{\mu\nu} \partial_\mu T \partial_\nu T}$$

Beyond the Veil: Uncovering the Truth About UFOs

where:

$V(T) = \tau_{\text{brane}} / \cosh(T/\sqrt{2})$ is the tachyon potential, fixed by Sen's boundary conditions:

- $V(0) = \tau_{\text{brane}}$ (the D-brane tension at the unstable maximum where ordinary matter is anchored)
- $V(T) \rightarrow 0$ as $T \rightarrow \infty$ (the tachyon vacuum, corresponding to the closed-string state identified in A2.6.1 as the “loop closure” step)

$$\eta^{\mu\nu} = \text{diag}(-1, +1, +1, +1)$$

is the Minkowski metric for the brane-world, and the field $T(t)$ is **real-valued** throughout the transition.

For spatially homogeneous configurations (the relevant case for a macroscopic object undergoing conversion, $T = T(t)$), the Euler–Lagrange variation yields the equation of motion:

$$\frac{d}{dt} \left[\frac{V(T)\dot{T}}{\sqrt{1 - \dot{T}^2}} \right] + \frac{V'(T)}{\sqrt{1 - \dot{T}^2}} = 0$$

For the canonical potential $V(T) = \tau_{\text{brane}} / \cosh(T/\sqrt{2})$, this admits the closed-form **rolling solution**:

$$T(t) = \sqrt{2} \cdot \ln \left[\cosh \left(\frac{t}{\tau_{\text{core}}} \right) \right]$$

with:

$$\tau_{\text{core}} = 1/\sqrt{|V''(0)|}$$

— the natural timescale set by the curvature of the potential at the unstable maximum, fixed by the D-brane tension (not a free parameter).

A2.4.3.3. The Chronology of Transmutation: From Sen's Rolling to the Revoredo Inversion Condition

The rolling solution (A2.4.3-iii) reproduces the stages of transmutation, but each stage now follows from the potential

Beyond the Veil: Uncovering the Truth About UFOs

$V(T)$ and its derivatives, rather than from a postulated activation profile:

Stage 1: Baryonic Ground State ($t \rightarrow -\infty$, $T = 0$)

The field sits at the unstable maximum of $V(T)$, where $V''(0) < 0$. The local effective mass-squared of small fluctuations is:

$$m_{\text{eff}}^2(t) = V''(T(t))$$

This is negative from the outset in the sense required for instability, while the field itself remains real throughout. The engine's Calabi-Yau torsion field T_{abc} (A2.6.2) selects this unstable extremum as the operating point, preventing the D-brane matter from settling into the stable vacuum it would otherwise occupy. It is worth clarifying that $m_{\text{eff}}^2 = V''(T)$ measures the stability of the tachyonic field T , not the mass of the ship or of its particles.

Stage 2: Orthogonal Rotation Phase ($0 < t < t_{\text{jump}}$)

The field $T(t)$ rolls smoothly away from $T = 0$. The effective mass-squared $m_{\text{eff}}^2(t) = V''(T(t))$ evolves as:

$$m_{\text{eff}}^2(t) = -\frac{\tau_{\text{brane}}}{2} \cdot \text{sech}^2\left(\frac{t}{\tau_{\text{core}}}\right) \cdot \tanh\left(\frac{t}{\tau_{\text{core}}}\right) \cdot \frac{1}{\cosh\left(\frac{T}{\sqrt{2}}\right)}$$

This is a **real-valued** function that crosses zero at the inflection point $T = T_{\text{inflection}}$ (where $V''(T) = 0$) and, once past the inflection, is positive and tends toward zero as the field approaches the tachyonic vacuum ($T \rightarrow \infty$): its instability is exhausted. The expression above illustrates the time variation; the exact definition is $m_{\text{eff}}^2 = V''(T(t))$, that is, $m_{\text{eff}}^2(T) = (\tau_{\text{brane}}/2) \cdot \text{sech}(T/\sqrt{2}) \cdot [\tanh^2(T/\sqrt{2}) - \text{sech}^2(T/\sqrt{2})]$, which equals $-\tau_{\text{brane}}/2$ at $T = 0$, vanishes at $T = \sqrt{2} \cdot \ln(1+\sqrt{2}) \approx 1.25$, and is positive from there on. The imaginary mass $m_0 = i\mu$ belongs to a different description of the same state: that of particles in the saturated regime (A2.2.1 and A2.4.3.4). The string endpoints undergo the Dirichlet→Neumann transition of A2.6.2 continuously, without the mass ever leaving the real

Beyond the Veil: Uncovering the Truth About UFOs
axis—resolving the non-Hermiticity concern that motivated earlier drafts.

Stage 3: The Singular Crossover ($t = t_{\text{critical}}$)

At the precise instant where $T(t) = T_{\text{inflection}}$, the effective mass-squared vanishes:

$$m_{\text{eff}}^2(t_{\text{critical}}) = 0$$

In standard 3D spacetime, a massless macro-object would trigger an immediate infinite-energy radiation catastrophe. However, because the endpoint detachment of the strings (A2.6.1) is occurring simultaneously via the torsion field T_{abc} , the field avoids the infinite-energy barrier. The energy-momentum tensor cancels its standard local confinement, preparing the topological transition.

Stage 4: Tachyonic Saturation ($t \geq t_{\text{jump}}$)

As $t \rightarrow +\infty$, $T \rightarrow \infty$, and $V(T) \rightarrow 0$. In this limit Sen's own analysis shows that the field's energy density approaches a nonzero constant while its pressure vanishes:

$$\rho_T \rightarrow \tau_{\text{brane}}, \quad p_T \rightarrow 0$$

This is the regime Sen names “**tachyon matter**” — a pressureless, geodesically moving distribution of closed-string excitations. This matches the closed-string, brane-independent character already assigned to the ship in tachyonic cruise (A2.4.1).

The Revoredo Inversion Condition (RIC) in Sen's Formalism

The Revoredo Engine must **actively sustain** the field within the unstable zone of the potential (from $T = 0$ to the inflection point) through the nonlocal energy compensation of the Revoredo-EPR Network, preventing it from “falling” to the real vacuum. This is achieved by the Phase Memory Tensor $M_{\mu\nu}$ (A2.6.3) acting as a boundary condition that enforces:

$$V''(T) < 0 \quad \text{and} \quad \partial_t(V(T)) > 0$$

This is the formal record of the Revoredo Inversion Condition: the engine maintains the field in a state of **inhibited tachyon condensation**, exactly at the boundary between ordinary matter and tachyonic matter. While the engine sustains it, the field $T(t)$ does not continue rolling to infinity: it is **held within the unstable zone, up to its edge at the inflection point**, by the active restoration term of the dynamic PMT. When the engine releases the field, it crosses the inflection and reaches tachyonic saturation (Stage 4). During cruise, the same active control holds the field's rate of change constant (φ , A2.8.6), with the help of the Time Crystals. The unstable maximum (the start), the path, and the inflection point (the edge) are thus distinct positions within a single zone of operation. The RIC is a postulate of the model.

A2.4.3.4. The Formal Bridge to PT-Symmetric Quantum Mechanics

The rolling solution (A2.4.3-iii) describes the transition from $m^2 > 0$ to $m^2 < 0$ in the **classical field limit**. However, the ship and its crew are quantum systems. The dynamic equation for the quantum state $|\Psi(t)\rangle$ during the rolling transition is governed by the Hamiltonian that emerges from Sen's action:

$$H_T = \int d^3x \sqrt{1 - \dot{T}^2} V(T) + (\text{momentum terms})$$

This Hamiltonian is **Hermitian** in the standard Dirac inner product, since $T(t)$ is real. The evolution of the density matrix $\varrho(t) = |\Psi(t)\rangle\langle\Psi(t)|$ is therefore unitary:

$$\partial_t \rho = -i[H_T, \rho]$$

However, once the system crosses the inflection point and enters the tachyonic saturation regime (Stage 4), the completed tachyonic state has $m_0 = i\mu$ (A2.2.1). The single-particle Hamiltonian for the tachyonic field is:

$$H_R = \frac{p^2}{2i\mu} + V_{CY}(x)$$

This is manifestly **non-Hermitian** in the standard Dirac inner product. The evolution is no longer unitary in the conventional sense. This is precisely the regime where PT-symmetric quantum mechanics becomes necessary.

A2.5. String Theory as Conceptual Underpinning

The original string theory (bosonic string theory) predicted exactly the scenario of the Revoredo Model: the first string to appear in the mathematics was a string vibrating with imaginary mass—that is, a tachyon. This was not discarded as an error, but interpreted as a fundamental state of the system.

A2.5.1. Particles as Vibration Modes

In string theory, elementary particles are not solid spheres but "musical notes" on a vibrating one-dimensional string. Changing the vibration mode (the harmonic) is equivalent to changing the particle. Mass, charge, and spin are properties of the vibration mode, not intrinsic properties of a point object.

A2.5.2. The Fundamental State of Hyperspace

In hyperspace (which in string theory requires 10 or 11 dimensions), space is not empty but filled with a fabric of strings in their fundamental state. If it were possible to control hyperspace, the key would not be to move a ship through it, but to alter the vibration mode of the fabric itself.

- **The tachyonic mode:** a string vibrating in the imaginary-mass mode alters the geometry of hyperspace, opening "shortcuts" or creating a phase in which traditional distances cease to exist as an operative concept.
- **The matter mode:** to re-enter the three-dimensional universe, the vibration is "retuned," reducing its energy to

Beyond the Veil: Uncovering the Truth About UFOs

transform back into electrons and quarks with real positive mass.

A2.5.3. The Geometry of Hidden Dimensions: Calabi-Yau Spaces

The additional dimensions of hyperspace are curled into complex geometric shapes called Calabi-Yau spaces. The precise way these dimensions are twisted determines how strings vibrate. If a technology capable of locally altering the geometry of those hidden dimensions existed, the strings composing an object would instantaneously change their vibration mode. In string theory (10 dimensions), the 6 extra dimensions are compactified on Calabi-Yau manifolds, and in M-theory (11 dimensions) the 7 extra ones are described by manifolds of G_2 holonomy. In this chapter, "Calabi-Yau space" designates, in a broad sense, the geometry of the compactified dimensions, and the model is formulated in terms of that geometry, whatever its exact description may be.

This is the central mechanism of the Revored Engine: changing the Calabi-Yau geometry in the immediate vicinity of the object to force its strings to adopt the tachyonic vibration mode.

A2.5.4. Imaginary Numbers as the Native Mathematics of Hyperspace

In the complex plane, the real and imaginary axes are orthogonal (perpendicular). If hyperspace exists at a "different angle" from our three-dimensional reality, the mathematics governing that direction could well be the mathematics of imaginary numbers. What to an observer on the real axis is an impossible or unstable mass would, on the hyperspace axis, be the standard and stable vibration of that environment.

A2.6. The Revoredo Tension Engine: Operation, Identity Preservation, and the Phase Memory Tensor Formalism

A2.6.1. Brana Resonance Modulation

In extensions of string theory (M-Theory), our three-dimensional universe could be a structure called a D-brane. The strings forming normal matter are anchored to this brane at their endpoints (open strings). The Revoredo Engine operates through three sub-processes:

- **Endpoint detachment:** a massive amount of Planck-scale localized quantum energy is injected to peel the string endpoints off the three-dimensional brane.
- **Loop closure:** upon detachment, open strings close on themselves, forming closed strings. Closed strings are not trapped in our brane and can move freely through hyperspace (the bulk).
- **Reverse retuning:** to return, the process reverses: the engine applies reverse tension to the strings, forcing them to open and re-anchor to the three-dimensional brane, instantaneously restoring positive mass and subluminal velocity, once the finite reconversion threshold has been reached (A2.8.5).

A2.6.2. The Catalyst: Tensorial Tension Fields

Modifying the vibration mode of a string requires more than pushing the macroscopic object; the underlying spacetime tension must be altered at the Planck scale (10^{-35} meters). The device generates a geometric deformation field that locally alters the compactified Calabi-Yau dimensions. By contracting or stretching these hidden dimensions, the string is forced to vibrate in a different harmonic, transitioning from positive mass to imaginary squared mass ($m^2 < 0$).

The geometric deformation field is formally a torsion field in Calabi-Yau space:

$$T_{abc} = \partial_a g_{bc}^{\text{CY}} - \partial_b g_{ac}^{\text{CY}}$$

where g_{ab}^{CY} is the metric of the compactified dimensions. The boundary conditions of open strings are sensitive to this geometry: a sufficiently intense torsion field in the extra dimensions inverts the Dirichlet boundary condition (endpoint fixed to the brane) to a Neumann boundary condition (free endpoint, zero momentum). The Neumann condition describes free endpoints: the inversion from Dirichlet to Neumann frees the endpoints from their anchoring, and the “loop closure” (A2.6.1) joins them to form a closed string, whose vibration modes include the bosonic tachyonic state of hyperspace. The binding energy released during detachment does not disappear: it converts into vibrational energy of the resulting closed string, configuring exactly the imaginary-mass state the model requires.

A2.6.3. The Phase Memory Tensor: Preserving Identity During the Transition

The Revoredo Engine must not only induce the mass-phase conversion, but must fully preserve the object's configuration information throughout the transition. If the strings forming a ship change their harmonic, how are the molecular structure, neuronal patterns, and memories of the crew preserved? In string theory, the vibration mode determines all quantum properties of the particle. A phase conversion without information preservation would be equivalent to destruction and recreation of the object—not transportation.

To resolve this problem, the Revoredo Engine incorporates an additional component: the Phase Memory Tensor (PMT). What follows is its derivation from first principles.

Stage 1 — The Object's Quantum State

Before the Revoredo Engine activates, the ship and all its contents form a quantum system of N strings. Their joint state is a pure state in the tensor-product Hilbert space:

$$|\Psi\rangle \in H^{\otimes N}$$

Beyond the Veil: Uncovering the Truth About UFOs

where each subspace H_i corresponds to an individual string with its quantum numbers: mass, spin, charge, isospin. This state is not a simple product of individual states. The strings forming an atom, a molecule, a neuron, are entangled in specific ways. That entanglement network is precisely what constitutes the structural identity of the object: a ship is not merely a list of strings, but a particular relational configuration of correlated strings.

Stage 2 — The Density Matrix

The state $|\Psi\rangle$ is encoded as the standard density matrix:

$$\rho = |\Psi\rangle\langle\Psi|$$

This is a Hermitian, positive semidefinite, unit-trace matrix. It contains all the information of the system: not only the expectation values of observables, but all quantum coherences—the interferences between distinct states that encode superposition and entanglement.

The relational correlator between the string pair (i, j) is extracted via:

$$C_{ij} = \text{Tr}[\rho \cdot (\sigma_i \otimes \sigma_j)]$$

where σ are the Pauli operators generalized to the string state space. The $N \times N$ matrix C_{ij} is the relational fingerprint of the object: a complete map of which strings are correlated with which others, and to what degree. If two distinct objects had exactly the same atoms but different neuronal configurations or different crystal structures, their C_{ij} matrices would differ. This is the level at which identity is encoded.

Why the Density Matrix Alone Is Insufficient

The density matrix ρ and the correlator C_{ij} are objects in Hilbert space, without μ - ν indices. They cannot appear directly in Einstein's equation, which operates in spacetime with covariant tensorial indices. A mathematical bridge is needed to project the quantum identity information into curved spacetime without losing coherences. That bridge is the Wigner-Weyl transform, followed by immersion via vierbein.

Beyond the Veil: Uncovering the Truth About UFOs

Stage 3 — The Wigner Function and Phase-Space Projection

The standard bridge between quantum mechanics and classical geometry is the Wigner-Weyl transform. The Wigner function projects the density matrix onto phase space (position, momentum):

$$W(x, p) = (1/\pi\hbar) \cdot \int \langle x+y | \rho | x-y \rangle \cdot e^{2ipy/\hbar} dy$$

This function has a fundamental property: it is not a classical probability distribution (it can take negative values, encoding the quantum nature of the state), but integrates correctly to give marginal probabilities in both position and momentum. It is a quasi-probabilistic representation of the quantum state in phase space, with the correct dimension to be integrated over a manifold.

The relational correlator becomes a position function:

$$C_{ab}(x) = \text{Tr}[\rho(x) \cdot (\sigma_a \otimes \sigma_b)]$$

where $\rho(x)$ is the local density of the density matrix at spacetime point x , obtained by integrating W over momenta. Indices a, b are local Lorentz frame (tetrad) indices, not yet curved-spacetime indices.

Stage 4 — Immersion in Curved Spacetime: Definition of the PMT

To pass from the local (a, b) Lorentz frame indices to the covariant (μ, ν) curved-spacetime indices, the vierbein e_μ^a is used—the standard object that relates both frames in general relativity. The Phase Memory Tensor is then defined as:

$$M_{\mu\nu} = \kappa \cdot e_\mu^a \cdot e_\nu^b \cdot \int W(x, p) \cdot C_{ab}(x) \cdot u_\alpha u_\beta g^{\alpha\beta} d^4x$$

The components of this expression have specific functions:

- $\kappa \sim 1/P^2 / \hbar$: the coupling constant, suppressed by the Planck scale. It ensures $M_{\mu\nu}$ is small compared to $T_{\mu\nu}^{\text{matter}}$ in the normal regime, and becomes dominant only when the operator Φ_R is active during the phase transition.

Beyond the Veil: Uncovering the Truth About UFOs

- **e_μ^a (vierbein):** the geometric object that maps the local Lorentz frame to curved spacetime. In flat spacetime it reduces to the Kronecker delta; in curved spacetime it encodes how the local frame is deformed by the geometry.
- **$W(\mathbf{x}, \mathbf{p})$:** the Wigner function weights the contribution of each phase-space region according to its quantum density. It supplies the quantum content of the identity state.
- **$C_{ab}(\mathbf{x})$:** the local relational correlator. It supplies the information about which strings are correlated at each point in the object—its identity structure.
- **$u_\alpha u_\beta g^{\alpha\beta}$:** is a scalar: the normalization of the four-velocity u of the comoving observer (it equals -1 with signature $(-, +, +, +)$ and $c = 1$). It fixes sign and scale; the tensor structure of $M_{\mu\nu}$ is supplied by the vierbeins and the correlator $C_{ab}(\mathbf{x})$.

The result is a symmetric, covariant rank-2 tensor that carries the complete quantum identity of the object as a field over spacetime. The derivation chain is:

$\varrho = |\Psi\rangle\langle\Psi| \rightarrow$ **complete quantum state (Hilbert space)**

$\downarrow$ Wigner-Weyl

$W(\mathbf{x}, \mathbf{p}) \rightarrow$ phase-space representation (semiclassical)

$\downarrow$ partial trace + correlator

$C_{ab}(\mathbf{x}) \rightarrow$ local relational correlations (local frame)

$\downarrow$ vierbein e_μ^a

$M_{\mu\nu} \rightarrow$ covariant tensor in curved spacetime

The PMT Conservation Law

For $M_{\mu\nu}$ to be physically consistent within the Revored field equation, it must satisfy the zero covariant-divergence condition—for the same reason $T_{\mu\nu}$ does in general relativity (consequence of the Bianchi identities):

$$\nabla^\mu M_{\mu\nu} = 0$$

Beyond the Veil: Uncovering the Truth About UFOs

This equation is the condition that the identity of the object is conserved along the hyperspace trajectory. It is the formal equivalent of saying: what enters hyperspace is exactly what exits. If this condition is violated, the object may emerge with a different identity—equivalent to destruction and recreation.

The Dynamic PMT with Evolution Equation

A statically formulated PMT would be fixed at the moment of conversion and remain unaltered. This would be physically naive. During the hyperspace transit, the tachyonic field continuously perturbs the state of the strings. A static PMT would irreversibly degrade under hyperspace dynamics.

An evolution equation for $M_{\mu\nu}$ along the affine parameter σ of the geodesic is therefore required:

$$\partial_{\sigma} M_{\mu\nu} = -i[H_{\text{CY}}, M_{\mu\nu}] - \gamma(M_{\mu\nu} - M_{\mu\nu}^{\text{target}}) + J_{\mu\nu}^{\text{EPR}}$$

Each term has a specific and irreplaceable function:

- $-i[H_{\text{CY}}, M_{\mu\nu}]$ (**unitary evolution**): the commutator with the Calabi-Yau Hamiltonian. Describes the unitary evolution of $M_{\mu\nu}$ under the local hyperspace geometry. It is the analog of the von Neumann equation for density-matrix evolution ($\partial_t \rho = -i[H, \rho]/\hbar$), translated to the curved spacetime of hyperspace. This term is unavoidable: any quantum tensor evolves under the Hamiltonian of the system in which it finds itself.
- $-\gamma(M_{\mu\nu} - M_{\mu\nu}^{\text{target}})$ (**active restoration**): a restoration term with rate γ . $M_{\mu\nu}^{\text{target}}$ is the reference state fixed at the moment of conversion, stored in the Quantum Time Crystals. If hyperspace perturbation pushes $M_{\mu\nu}$ away from its initial state, this term drives it back with a force proportional to the deviation. It is formally identical to a Lindblad dissipation term, but operating in reverse: instead of destroying coherence, it actively restores it. The rate γ must exceed the tachyonic

Beyond the Veil: Uncovering the Truth About UFOs

perturbation rate—a condition the Time Crystals guarantee.

- **$J_{\mu\nu}^{\text{EPR}}$ (entangled correction):** the entangled correlation current of the Revoredo-EPR Network. This term continuously provides a shared reference about the target state through the Dyson Sphere. Through the ER=EPR channel, quantum correlations between the entangled counterpart of the PMT in the Sphere and the tensor in the ship are used to correct deviations in real time.

The explicit form of the entangled correction term is:

$$J_{\mu\nu}^{\text{EPR}} = \lambda \cdot \text{Tr}_{\text{EPR}} \left[\Omega_{\mu\alpha\nu\beta} \cdot \rho_{\text{EPR}}^{\alpha\beta} \right]$$

where $\Omega_{\mu\alpha\nu\beta}$ is the Calabi-Yau phase curvature tensor (already present in the geodesic equations), $\rho_{\text{EPR}}^{\alpha\beta}$ is the density matrix of the entanglement channel, and λ is the EPR coupling constant. This expression states that the correction established from the entangled counterpart in the Dyson Sphere is filtered through the local hyperspace geometry before being applied to the memory tensor. The quantum state of the ship is not cloned (which would contradict the no-cloning theorem): what is recorded is the pattern of correlations (the target PMT), and the counterpart in the Sphere is the other end of one and the same entanglement, a relational structure seen from two places. Aboard the ship, the reference anchor of that pattern is the Time Crystals. That correlations suffice to define identity is a postulate of the model.

The Complete PMT System

The Phase Memory Tensor is defined by the following system of equations:

Definition:

$$M_{\mu\nu} = \kappa \cdot e_{\mu}^a \cdot e_{\nu}^b \cdot \int W(x, p) \cdot C_{ab}(x) \cdot u_{\alpha} u_{\beta} g^{\alpha\beta} d^4x$$

Evolution:

$$\partial_{\sigma} M_{\mu\nu} = -i[H_{\text{CY}}, M_{\mu\nu}] - \gamma(M_{\mu\nu} - M_{\mu\nu}^{\text{target}}) + J_{\mu\nu}^{\text{EPR}}$$

Beyond the Veil: Uncovering the Truth About UFOs

Conservation:

$$\nabla^\mu M_{\mu\nu} = 0$$

Fidelity:

$$|| M_{\mu\nu}(\sigma) - M_{\mu\nu}^{\text{target}} || < \epsilon_{\text{critical}} \text{ [condition prior to reconversion]}$$

This system closes the PMT formalism in a mathematically consistent manner. Its level of speculation is the same as the rest of the model: it rests on real mathematical objects (density matrix, Wigner function, vierbein, von Neumann equation, Lindblad formalisms) coherently extended toward an experimentally unverified regime—exactly as string theory does with its own extensions.

A2.6.4. Stability of the Phase Memory Tensor Under Non-Hermitian Evolution: A PT-Symmetric Formulation Derived from Sen's Action

A2.6.4.1. The Problem: Non-Hermiticity in the Tachyonic Phase

The rolling tachyon transition (A2.4.3) keeps the field $T(t)$ real, and the associated Hamiltonian H_T is Hermitian. However, once the ship has fully entered the tachyonic saturation regime (Stage 4), the effective rest mass is $m_0 = i\mu$ (A2.2.1). The Hamiltonian for a single string in the tachyonic state is:

$$H_R = \frac{p^2}{2i\mu} + V_{CY}(x)$$

where $V_{CY}(x)$ is the Calabi-Yau torsion potential that governs the interaction with the compactified dimensions. This Hamiltonian is non-Hermitian in the ordinary Dirac inner product ($H_R \neq H_R^\dagger$).

Without a correction, the density matrix ρ would evolve non-unitarily, leading to decoherence and the destruction of the relational fingerprint C_{ij} (A2.6.3) that encodes the object's identity. The Phase Memory Tensor $M_{\mu\nu}$ would degrade, and

Beyond the Veil: Uncovering the Truth About UFOs

the ship would emerge from hyperspace as a different object—or not at all.

A2.6.4.2. The PT-Symmetric Resolution: Unitarity in the CPT Norm

Non-Hermitian Hamiltonians of exactly this kind—complex mass or complex potential terms that nonetheless respect combined parity-time (PT) symmetry—were shown by Bender and Boettcher (1998) to possess entirely real energy spectra over a broad parameter range. Bender, Brody, and Jones (2002) subsequently proved that such Hamiltonians generate fully unitary time evolution once the physical inner product is redefined using the **CPT norm**, built from a positive-definite metric operator $\eta_R = e^{(-Q_R)}$.

The construction is as follows:

1. Define the PT operator: Let P be the parity operator (spatial inversion in the compactified Calabi-Yau dimensions) and T the time-reversal operator. The PT operator acts on the field and potential as:

$$\text{PT: } x \rightarrow -x, \quad p \rightarrow -p, \quad i \rightarrow -i$$

2. Impose PT symmetry: The Hamiltonian H_R satisfies:

$$[H_R, \text{PT}] = 0$$

In terms of the potential, this requires:

$$V_{\text{CY}}(x) = V_{\text{CY}}^*(-x)$$

This is a condition the model can plausibly assign to the torsion field T_{abc} of A2.6.2, since that field is already antisymmetric in its indices and lives in the compactified dimensions. A clarification on the kinetic term: with the usual action of T, which leaves mass invariant, the term $p^2/(2i\mu)$ would change sign under $i \rightarrow -i$ and break PT symmetry. The model postulates that, in hyperspace, T additionally inverts the geometric chirality of the Calabi-Yau dimensions, so that the effective mass parameter changes sign under time reversal ($\mu \rightarrow$

Beyond the Veil: Uncovering the Truth About UFOs

$-\mu$). Thus $p^2/(2(-i)(-\mu)) = p^2/(2i\mu)$ remains invariant and $[H_R, PT] = 0$ is consistent. Whether the spectrum turns out real also depends on H_R remaining in the unbroken PT phase (A2.6.4.4).

3. Construct the metric operator: There exists a positive-definite operator $\eta_R = e^{(-Q_R)}$ such that:

$$h_R = \eta_R^{1/2} H_R \eta_R^{-1/2}$$

is an ordinary Hermitian Hamiltonian, unitarily equivalent to H_R . The spectrum of H_R is real if and only if H_R is in the **unbroken PT-symmetric phase**.

4. Redefine the inner product: The physically meaningful inner product is the **CPT inner product**:

$$\langle \varphi | \psi \rangle_{\text{CPT}} = \langle \varphi | \eta_R | \psi \rangle$$

Under this inner product, H_R is Hermitian:

$$\langle H_R \varphi | \psi \rangle_{\text{CPT}} = \langle \varphi | H_R \psi \rangle_{\text{CPT}}$$

and the evolution is unitary.

A2.6.4.3. The PT-Symmetric Phase Memory Tensor

The Phase Memory Tensor (A2.6.3) must be reformulated in the PT-symmetric framework. The density matrix $\varrho = |\Psi\rangle\langle\Psi|$ is redefined as the **CPT density matrix**:

$$\rho_{\text{PT}} = \eta_R^{1/2} \varrho \eta_R^{-1/2}$$

The relational correlator C_{ij} of A2.6.3 is computed using the CPT trace:

$$C_{ij} = \text{Tr}_{\text{CPT}}[\rho_{\text{PT}} \cdot (\sigma_i \otimes \sigma_j)]$$

where:

$$\text{Tr}_{\text{CPT}}[\mathcal{O}] = \text{Tr}[\eta_R \mathcal{O}]$$

The Phase Memory Tensor becomes:

$$M_{\mu\nu} = \kappa \cdot e_\mu^a e_\nu^b \cdot \int W(x, p) \cdot C_{ab}^{\text{PT}}(x) \cdot u_\alpha u_\beta g^{\alpha\beta} d^4x$$

where $C_{ab}^{\text{PT}}(x)$ is the PT-symmetric relational correlator.

Beyond the Veil: Uncovering the Truth About UFOs

The dynamic evolution equation for $M_{\mu\nu}$ is unchanged in form:

$$\partial_{\sigma} M_{\mu\nu} = -i[H_{CY}, M_{\mu\nu}] - \gamma(M_{\mu\nu} - M_{\mu\nu}^{\text{target}}) + J_{\mu\nu}^{\text{EPR}}$$

But the commutator term is now understood to generate exactly unitary evolution in the CPT norm, with no residual decoherence to compensate. The restoration term γ and the EPR correction $J_{\mu\nu}^{\text{EPR}}$ keep their original roles—active restoration against external hyperspace perturbation—rather than being pressed into service against an artifact of the Hamiltonian's own non-Hermiticity.

A2.6.4.4. The PT-Symmetry Preservation Condition (PSC)

The single, sharp requirement replacing any arbitrary inequality is that $H_R(t)$ remains in the **unbroken PT-symmetric phase** throughout the transmutation window and the hyperspace cruise:

$$\text{Spec}[H_R(t)] \subset \mathbb{R} \quad \forall t \in [0, t_{\text{return}}]$$

If this condition fails—if H_R undergoes spontaneous PT-symmetry breaking—pairs of its eigenvalues become complex conjugates of one another:

$$E_n \rightarrow E_n \pm i\Gamma_n \quad \text{with } \Gamma_n > 0$$

The corresponding PMT eigenmodes lose their physical (real, positive) norm under η_R . The relational fingerprint C_{ij} ceases to be well-defined, and identity preservation genuinely fails.

This gives the model a single, well-defined failure criterion: **PT-symmetry breaking**, the same criterion that governs the well-studied laboratory realizations of these Hamiltonians in optics and open quantum systems.

A2.6.4.5. The PT-Symmetric Bridge to Sen's Rolling Tachyon²

The unification is completed by showing that the PT-symmetric Hamiltonian H_R of A2.6.4-i is the **quantum counterpart** of the classical rolling tachyon action S_T of A2.4.3-i. This is achieved through the following steps:

1. Canonical quantization of Sen's action: Promote the classical field $T(t)$ to a quantum operator $\hat{T}$. The resulting quantum Hamiltonian is:

$$\hat{H}_T = \int d^3x \sqrt{1 - \hat{T}^2} V(\hat{T})$$

2. PT-symmetric extension: In the tachyonic saturation limit $T \rightarrow \infty$, $V(T) \rightarrow 0$, the Hamiltonian reduces to:

$$\hat{H}_T \xrightarrow{T \rightarrow \infty} \frac{\hat{p}^2}{2i\mu} + V_{CY}(\hat{x})$$

which is exactly H_R of A2.6.4-i. The PT symmetry of H_R is a **consequence** of the Calabi-Yau geometry, not an external imposition.

3. Unified action: The complete Revoredo action, covering both the rolling transition and the PT-symmetric tachyonic phase, is:

²The unification between Sen's rolling tachyon formalism and PT-symmetric quantum mechanics is presented here as a **working hypothesis**, grounded in the structural correspondence between both frameworks. A full derivation of the unified action (Eq. A2.6.4-xvi) would require a detailed canonical quantization of the tachyon field in the high-energy limit, including the explicit emergence of the $1/(2i\mu)$ kinetic term and the coupling to $V_{CY}(x)$. These steps are not shown in full; their rigorous completion is left for future work. Nonetheless, the plausibility of the connection rests on the shared mathematical architecture (hyperbolic-cosh potential and PT symmetry), and the resulting unified framework is internally consistent within the scope of this model.

$$S_{\text{Revoredo}} = \int d^{10}x \sqrt{-g} \left[\frac{1}{2\kappa^2} R - V(T) \sqrt{1 - \eta^{\mu\nu} \partial_\mu T \partial_\nu T} + \mathcal{L}_{\text{PT}} \right]$$

where $\mathcal{L}_{\text{PT}}$ contains the PT-symmetric terms that become dominant in the tachyonic regime.

A2.6.4.6. The Final Fidelity Condition

The condition for safe reconversion from the tachyonic state to ordinary matter is the simultaneous satisfaction of:

1. PT-symmetry preservation: The Hamiltonian H_R must remain in the unbroken phase throughout the journey:

$$\text{Spec}[H_R(t)] \subset \mathbb{R} \quad \forall t \in [0, t_{\text{return}}]$$

2. PMT fidelity threshold: The Phase Memory Tensor must remain within the critical fidelity envelope:

$$\|M_{\mu\nu}(\sigma) - M_{\mu\nu}^{\text{target}}\| < \epsilon_{\text{critical}}$$

3. Hyperspace Positioning System confirmation: The HPS readings (A2.12.4) must confirm the ship is in the quantum vicinity of the intended destination.

If and only if all three conditions are met, the Revoredo Engine can initiate the reverse retuning (A2.6.1), restoring the ship to ordinary baryonic matter with its identity intact.

A2.7. The Energy Source: Dyson Sphere and Zero-Point Energy

A2.7.1. Zero-Point Energy

Zero-point energy (ZPE) is the lowest possible energy state of a quantum physical system: the energy of the vacuum itself. Due to Heisenberg's Uncertainty Principle, the vacuum is never truly empty, but seethes with quantum fluctuations (virtual particles that constantly appear and disappear). To extract useful energy from this quantum sea, the system induces an effect analogous to the Casimir Effect at macroscopic scale: by geometrically constraining the vacuum's vibration modes in

Beyond the Veil: Uncovering the Truth About UFOs

specific regions, a quantum pressure differential is generated that can be converted into work or net energy.

A2.7.2. The Vacuum Catastrophe: The 120-Order-of-Magnitude Discrepancy

Physics confronts here one of its deepest contradictions. Two radically different calculations of vacuum energy density exist:

- **Quantum calculation (Planck density):** quantum equations predict a density of $\sim 10^{93}$ g/cm³. A single cubic centimeter of empty space would contain more mass-energy than the entire observable universe ($\sim 10^{56}$ grams).
- **Cosmological measurement (dark energy):** the value measured by telescopes is $\sim 10^{-29}$ g/cm³, equivalent to the mass of about 5 hydrogen atoms per cubic meter of space.

The discrepancy between these values is 120 orders of magnitude: the largest prediction error in modern physics. This gap is itself evidence that our understanding of the quantum vacuum is incomplete, making that unknown territory the domain in which the Revoredo Model situates its energy hypothesis.

A2.7.3. The Revoredo Model's Window

If quantum theory is right, the vacuum is an infinite ocean of energy that is blocked or canceling itself in ways we do not yet understand. A civilization capable of building a quantum Dyson Sphere would not need to create energy; it would only need to develop the technology to locally unlock an infinitesimal fraction of that quantum potential. By extracting the latent energy of a volume the size of a grain of sand, it would have enough mass-energy to reach the Planck scale and alter the string vibrations of any spacecraft.

A2.7.4. The Channel: Nonlocal Energy Compensation via Quantum Entanglement (ER=EPR)

The ER=EPR conjecture (Maldacena & Susskind) proposes that quantum entanglement and wormholes are mathematically the same thing. Each entangled pair of particles is connected by a micro-wormhole at the Planck scale. If the Dyson Sphere's particles are massively entangled with equivalent particles in the ship, a direct quantum correlation channel is created. This channel does not send energy through space. The mechanism consists of obtaining energy from the vacuum at the ship's location and paying it back, automatically and regardless of distance, in the vacuum of the Dyson Sphere, through the entangled correlation of the Revoredo-EPR Network: the energy that is obtained is not the same as the energy that is paid, and entanglement does not carry energy or information. It is analogous to a bank transfer between countries: the money withdrawn from one bank is not the same money that was deposited in another. That this compensation occurs without any signal is a postulate of the model.

A2.7.5. The Quantum Dyson Sphere

In the Revoredo Model, the Dyson Sphere is conceived as a megastructure for quantum computation and manipulation, with three functions:

- ZPE extraction via macroscopic Casimir effect in circumstellar space.
- Phase stabilization: maintains stable quantum fields and coordinates the massive entanglement of particles.
- Energy routing: acts as the central node of the Revoredo-EPR Network, feeding the geometric torsion fields required for phase conversion in the ship.

From a thermodynamic standpoint, the Dyson Sphere functions not only as an energy source but also as an entropy sink. The local reduction of entropy required to maintain the ship's quantum state and structural identity during tachyonic transit is compensated by the massive energy dissipation and

Beyond the Veil: Uncovering the Truth About UFOs

entropy increase occurring in the stellar-scale infrastructure of the Dyson Sphere. The total entropy of the system — ship, quantum fields, and the Dyson Sphere environment — never decreases, satisfying the second law of thermodynamics. A quantitative balance of entropy production is left as a development for future work.

A2.8. The Required Energy Scale: Planck Energy and the Asymmetry of the Return Journey

A2.8.1. Planck Energy

Altering the vibration modes of strings and forcing phase conversion requires Planck Energy: the fundamental limit at which the fabric of spacetime breaks down and the current laws of physics cease to apply.

$$E_P = m_P \cdot c^2 \approx 1.956 \times 10^9 \text{ Joules per individual string}$$

Concentrating nearly two billion joules within the size of a single string (10^{-35} meters) represents an unimaginable energy density: roughly equivalent to the chemical energy of a full aviation fuel tank, compressed into a point trillions of times smaller than an atom.

A2.8.2. Comparison With Current Technology

The Large Hadron Collider (LHC) at CERN operates at ~ 13 TeV. The required Planck Energy is 1.22×10^{19} GeV. The gap is 15 orders of magnitude: an energy one trillion times greater than what our most advanced technology can currently generate. The Dyson Sphere, capturing the total output of a star ($\sim 3.8 \times 10^{26}$ watts), would have the power needed to generate these pulses continuously.

A2.8.3. The Intrinsic Tension of Strings

The reason for this extreme energy requirement is the inherent tension of strings. For strings to generate particles with the diminutive masses we observe, they must be incredibly rigid.

Beyond the Veil: Uncovering the Truth About UFOs

Their theoretical tension is inversely proportional to the square of their length:

$$\text{String tension} \approx 10^{43} \text{ Newtons}$$

To modify the vibration mode of an object with such a basal tension, the applied energy pulse must match or exceed that force—which takes us directly back to the Planck scale.

A2.8.4. The Energetic Asymmetry of the Return Journey

The natural velocity of the tachyonic state is infinity, and braking requires energy. This creates a fundamental energetic asymmetry between the outbound and return journeys. The initial jump requires Planck Energy for phase conversion; but the return also requires additional energy to reduce the basal tachyonic velocity to a safe threshold before the matter can be reconverted.

A tachyonic-state object attempting to reconvert directly while traveling at infinite velocity would face a mathematical inconsistency: reversion requires a well-defined real proper time τ , but at infinite velocity the affine parameter σ of hyperspace and the proper time τ of real space become completely decoupled.

A2.8.5. Phase Asymmetry Term in the Revored Phase-Inversion Metric

A phase asymmetry term is incorporated into the Revored Phase-Inversion Metric:

$$\Delta_E = \frac{E_{\text{return}}}{E_{\text{departure}}} > 1$$

This term represents the energy gradient required to reduce tachyonic velocity from infinity to the minimum safe reversion threshold ($v_{\text{min}} > c$ but finite), where the affine parameter σ can once again couple to the proper time τ . There, the engine reverts the phase to baryonic matter, which reappears in a subluminal regime: just as on the outbound trip, the barrier is not crossed but bypassed through a phase change.

Beyond the Veil: Uncovering the Truth About UFOs

The Revored Engine operates in three modes: Conversion Mode (initial jump, cost: Planck Energy); Cruise Mode (tachyonic navigation, cost: directional and orientation energy); and Tachyonic Deceleration Mode (velocity reduction prior to reconversion, cost: variable depending on cruise speed).

In terms of mission design, the Dyson Spheres at the point of origin must supply approximately twice the energy required for the pure jump, reserving half for return-journey energy support.

A2.8.6. Equation of State of the Tachyonic Field During Cruise

The thermodynamic state of the field during the cruise transit—the period between phase conversion and tachyonic deceleration prior to reconversion—can be formally described using the tachyonic matter formalism. For a scalar field with $m^2 < 0$ in the superluminal regime, the equation of state is:

$$w_R = \frac{P_R}{\rho_R}$$

where P_R is the effective pressure of the field and ρ_R is its effective energy density. The explicit expressions are:

$$\rho_R = \frac{-V(\varphi)}{\sqrt{1 - (\partial_t \varphi)^2}}$$

$$P_R = -V(\varphi) \cdot \sqrt{1 - (\partial_t \varphi)^2}$$

yielding $w_R = -(1 - (\partial_t \varphi)^2)$, which oscillates between -1 and 0 . Under the Revored Inversion Condition, the Engine stabilizes $\partial_t \varphi$ at a fixed cruise value φ_c , producing a rigid equation of state during transit:

$$w_R^{\text{cruise}} = -(1 - \varphi_c^2) = \text{constant}$$

This state is analogous to that of the cosmological constant ($w = -1$) and is formally treatable as a stable relativistic fluid

Beyond the Veil: Uncovering the Truth About UFOs

over the Riemannian manifold of the modified metric $g_{\mu\nu}^R$. Quantum Time Crystals act as absolute phase-reference quantum oscillators that prevent drift of the cruise state: if the field tends to deviate from φ_c , their spontaneous temporal periodicity applies a restoring tension equivalent to the $-\gamma$ term of the dynamic PMT. The condition for exiting the cruise state is simultaneous verification by the SPH of the quantum vicinity of the destination, and of the PMT fidelity threshold, before initiating tachyonic deceleration.

A2.9. Official Nomenclature of the Model

In recognition of the development of this theoretical hypothesis, the mechanism receives the following formal designation within the framework of advanced theoretical physics:

Revoredo Quantum Transition Effect (Revoredo Mechanism)

The key components of the system are formally defined as follows:

- **Revoredo Tension Engine:** the onboard device that channels Planck-scale energy to alter Calabi-Yau's hidden dimensions and force the phase conversion of the ship's strings, incorporating the Phase Memory Tensor to preserve the identity of the object.
- **Revoredo Phase-Inversion Metric:** the set of mathematical equations describing how a real-matter open string mutates its vibration harmonic into a closed tachyonic string with negative squared mass ($m^2 < 0$), including the energetic asymmetry term for the return journey.
- **Revoredo-EPR Distribution Network:** the quantum entanglement bridge powered by the Dyson Sphere that nonlocally compensates for the vacuum energy obtained by the ship regardless of its hyperspace coordinates, and

Beyond the Veil: Uncovering the Truth About UFOs

also serves as the infrastructure for the Hyperspace Positioning System.

A2.10. The Temporal Paradox of Revoredo Travel

A2.10.1. Inverse Time Dilation

When the Revoredo Mechanism activates, the crew's perception and flow of time inside hyperspace undergo a fundamental alteration imposed by the mathematics of extended special relativity:

- **Imaginary Time:** in our normal space ($v < c$), time passes more slowly for the traveler at higher speeds (standard time dilation). Upon crossing the barrier via the Revoredo Effect ($v > c$), the Lorentz factor becomes an imaginary number. Conventional time ceases to advance in the ordinary direction.
- **Protected Causality:** for crew in the tachyonic phase, the journey is not experienced as chaotic temporal displacement. With closed strings in hyperspace, the ship experiences a proper-time state orthogonal to ours—perpendicular to the three-dimensional temporal flow. Formally, the affine parameter σ parametrizing the hyperspace geodesic is mathematically independent of the real-space proper time τ during transit, which suppresses the conventional causal channels through which classical temporal paradoxes could propagate.
- **Apparent Instantaneous Effect:** upon re-entering real space via phase reversion, for the external universe the journey will have taken a time proportional to the distance in hyperspace. For the crew aboard, the trip will have been practically instantaneous: in the pure tachyonic state, resistance to standard temporal flow is zero.

A2.11. First Field Equation of the Revoredo Mechanism

A2.11.1. Tensorial Formulation

To formalize the hypothesis at the same level of speculative rigor as General Relativity or string theory, the fundamental field equation of the Revoredo Quantum Transition Effect is formulated. In Einstein's General Relativity, spacetime geometry is linked to matter through:

$$G_{\mu\nu} = \left(\frac{8\pi G}{c^4} \right) \cdot T_{\mu\nu}$$

For the Revoredo Mechanism, a quantum phase-transition operator Φ_R is introduced that acts directly on the Energy-Momentum Tensor ($T_{\mu\nu}$), forcing conventional matter to acquire the algebraic properties of a tachyon ($m^2 < 0$). The unified field equation of the Revoredo Model, including the Phase Memory Tensor, is:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \left(\frac{8\pi G}{c^4} \right) \cdot [T_{\mu\nu}^{\text{matter}} + \Phi_R \cdot T_{\mu\nu}^{\text{ZPE}} + M_{\mu\nu}^{\text{TMF}}]$$

where the energy tensor coupled to the Revoredo field is decomposed at the Planck scale as:

$$T_{\mu\nu}^{\text{Revoredo}} = (\rho_{\text{ZPE}} c^2 + T_{\text{string}}) \cdot u_\mu u_\nu + P_{\text{Casimir}} \cdot (g_{\mu\nu} + u_\mu u_\nu)$$

Analysis of the Tensor Components

- $T_{\mu\nu}^{\text{ZPE}}$: the Zero-Point Energy Tensor. Represents the immense latent energy density of the quantum vacuum ($\sim 10^{93}$ g/cm³).
- **Φ_R (Revoredo Phase Matrix Operator)**: a higher-order tensor derived from the additional Calabi-Yau dimensions. When the engine activates, it commutes from a real scalar state to a pure complex one: $\Phi_R \rightarrow i \cdot \chi$. This transition is the formal record of the Revoredo Inversion Condition: as the operator becomes complex, the scalar

Beyond the Veil: Uncovering the Truth About UFOs

field φ enters the regime of inhibited tachyon condensation, producing $T_{00} < 0$ locally as derived in section A2.4.2.

- **T_{string} (String Tension):** represents the fundamental rigidity of strings at the Planck scale ($\sim 10^{43}$ N). As the operator becomes complex, the effective tension inverts algebraically: $T_{\text{string}} \rightarrow -|T_{\text{string}}|$.
- **$M_{\mu\nu}^{\text{TMF}}$ (Phase Memory Tensor):** an auxiliary field derived from first principles (section A2.6.3) via the chain $\varrho \rightarrow W(x,p) \rightarrow C_{ab}(x) \rightarrow M_{\mu\nu}$. Preserves the relational configuration of strings during the transition, guaranteeing the structural integrity and identity of the object and its crew across the mass-regime change. Obeys its own evolution equation and conservation law.

The sign change in the energy tensor is the precise mathematical record of tachyon condensation. It causes the energy-momentum tensor to locally describe a pure imaginary mass, enabling the object to travel in superluminal trajectories ($v > c$) without violating local energy conservation.

A2.12. Geodesic Trajectory Equations and the Hyperspace Positioning System

A2.12.1. The Revoredo Modified Metric ($g_{\mu\nu}^R$)

The phase operator $\Phi_R = i\chi$ (where χ is the hyperspace coupling factor) alters the spacetime metric. The effective Revoredo metric is expressed as:

$$g_{\mu\nu}^R = g_{\mu\nu} - (1 + \chi^2) \cdot u_\mu u_\nu$$

where $g_{\mu\nu}$ is the background metric, u_μ is the four-velocity vector of the distortion bubble, and χ is the quantum transition parameter. When the engine is off $\chi = 0$, the metric reduces exactly to $g_{\mu\nu}$, recovering the standard background metric. The line element in the direction of travel (x-axis) transforms as:

Beyond the Veil: Uncovering the Truth About UFOs

$$ds^2 = -c^2 dt^2 + \chi^2(x) \cdot (dx - v_s \cdot dt)^2 + dy^2 + dz^2$$

where v_s is the propagation velocity of the quantum wave front generated by the Calabi-Yau graphene resonators. When $\chi^2 > 1$, the effective sign of the temporal component is locally inverted inside the bubble.

The Modified Christoffel Symbols

The trajectory is determined by the tidal forces encoded in the Christoffel symbols. For the modified metric $g_{\mu\nu}^R$, the Christoffel symbols are computed in the standard way from the metric and its derivatives:

$$\Gamma_{tt}^x = \frac{1}{2} \cdot \chi^2 \cdot v_s \cdot \left(\frac{\partial \chi}{\partial x} \right) + \left(\frac{8\pi G}{c^4} \right) \cdot (T_{string} \cdot u_t)$$

This term demonstrates that the spatial gradient of the Revored phase field acts as an artificial repulsive gravity ahead of the ship and attractive gravity behind it.

A2.12.2. The General Revored Geodesic Equation

The ship is parametrized with respect to a hyperspace affine parameter σ (not with respect to proper time τ , which becomes pure imaginary during the tachyonic transition):

$$\frac{d^2 x^\mu}{d\sigma^2} + \Gamma_{\alpha\beta}^\mu \cdot \left(\frac{dx^\alpha}{d\sigma} \right) \left(\frac{dx^\beta}{d\sigma} \right) = F_{\text{Topological}}^\mu$$

where the non-gravitational quantum-origin force vector is:

$$F_{\text{Topological}}^\mu = \left(\frac{q_R}{m_{\text{imaginary}}} \right) \cdot \Omega_v^\mu \cdot \left(\frac{dx^\nu}{d\sigma} \right)$$

Pilot Control Variables:

- $m_{\text{imaginary}} = \mathbf{i} \cdot m_0$: the effective mass of the ship after tachyon condensation. Being imaginary, it counteracts the infinite resistance terms of Special Relativity.

Beyond the Veil: Uncovering the Truth About UFOs

- **qR (Revoredo charge):** a macroscopic quantum property determined by the entanglement state of the Majorana fermions in the engine-core shielding.
- **Ω^μ_ν (Calabi-Yau Phase Curvature Tensor):** controls the direction of the thrust vector in the compactified extra dimensions. It is the system's direction and navigation parameter.

A2.12.3. The Revoredo-EPR Hyperspace Positioning System

The geodesic trajectory equations are mathematically solid, but they raise a critical navigation gap: in tachyonic hyperspace, with no three-dimensional brane as a reference, how does the ship orient its trajectory toward its destination? The Calabi-Yau tensor Ω controls direction but requires knowing the local hyperspace geometry at every point—equivalent to having a prior map of the bulk.

This problem is resolved within the model itself using the Revoredo-EPR Network as a positioning system. Entangled nodes in Dyson Spheres distributed throughout the galaxy or cluster act as reference beacons. The ship, being entangled with each of these nodes, measures the phase differences of the quantum correlation with each beacon. This reading does not imply a triangulation based on conventional spatial waves. The model postulates, drawing on Susskind's "Complexity = Volume" ($C=V$) conjecture, that the geometry of the bulk is structurally encoded in the entanglement network with the nodes of the Dyson Sphere, established in advance: the ship receives no signal at all, but rather reads that structure of correlations. Assuming a technology capable of extracting geometric information from strongly correlated states, the system derives its hyperspace coordinates by relying on this correspondence between purely quantum information and spatial geometry.

A2.12.4. Hyperspace Positioning System (HPS Revored-EPR)

Since entanglement has no effective distance in hyperspace (operating through ER=EPR micro-wormholes), phase variations encode relative position information in the bulk. The ship constructs its hyperspace coordinates in real time: an intergalactic quantum GPS whose precision increases with the number of available entangled nodes.

This system also resolves the problem of safe reconversion at the destination: the ship does not initiate reverse conversion until HPS readings confirm it is in the quantum vicinity of the intended destination.

The Revored-EPR Network thus fulfills a triple function: channel for nonlocal energy compensation with the Dyson Sphere, high-precision hyperspace navigation system, and continuous correction channel for the Phase Memory Tensor.

A2.13. Stable Quantum Materials for the Revored Engine

The Revored Engine cannot be built from ordinary baryonic matter. Any normal atom would disintegrate when subjected to Planck Energy Density. What is required are quantum materials with intrinsic topological engineering capable of interacting with hyperspace and sustaining massive EPR entanglement currents.

A2.13.1. Topological Majorana Fermion Condensates (Core Shielding)

Majorana fermions are particles that are their own antiparticles and exhibit non-Abelian statistical properties (anyons). In a topologically protected state, they are immune to quantum decoherence induced by the external environment.

- **Function:** to clad the engine core. Acts as an absolute quantum thermal superconductor.
- **Mechanism:** channels zero-point energy extracted by the Dyson Sphere and distributes it uniformly to the Planck injectors without dissipating it as destructive heat.

A2.13.2. Calabi-Yau Graphene Metamaterials (String Resonators)

Two- and three-dimensional structures arranged at the nanometric level that mimic the compactified geometry of hyperspace.

- **Function:** to physically tune the string vibration modes of the ship, and to generate the torsion field in the Calabi-Yau dimensions required for the Dirichlet-Neumann boundary-condition transition of the open strings.
- **Mechanism:** via manipulation of molecular quantum potential wells, they act as molecular "guitar picks." When an energy current passes through them, they resonate at the precise frequencies needed to alter the tensor Φ_R , mechanically forcing the phase change of surrounding baryonic matter to tachyonic matter.

A2.13.3. Quantum Time Crystals (Phase Stabilizers)

Structures whose patterns repeat not only in space, but also in time—perpetually, without consuming energy. Time crystals realized in the laboratory are periodically driven; the model assumes an idealized version, in the lowest-energy state, characteristic of a technology far beyond what exists today. This is an assumption of the model.

- **Function:** to maintain the ship stable in the imaginary-mass state during hyperspace travel, to stabilize the cruise-state value φ_c of the tachyonic field, and to prevent premature tachyon condensation (accidental reversion to real mass in the middle of hyperspace).
- **Mechanism:** the time crystal acts as an absolute quantum phase reference clock. It locks the quantum phase of the ship's strings in a specific vibration, breaking the field's natural tendency to decay toward the stable vacuum. It also stores the $M_{\mu\nu}^{\text{target}}$ state of the Phase Memory Tensor, acting as the physical repository of the object's identity during hyperspace transit.

A2.14. Comparison: Revoredo vs. Alcubierre for Kardashev Civilizations

A2.14.1. For Type II and Type III Civilizations

For a Type II civilization (capable of harnessing its star's total energy) or a Type III civilization (capable of controlling the energy of an entire galaxy), superluminal transport would not be a matter of choosing a single technology, but of applying the appropriate mechanism according to strategic need. Comparing the Alcubierre Metric with the Revoredo Tachyonic Conversion Mechanism, we see that an advanced civilization would have two radically different physical methodologies that complement each other.

Dimension	Revoredo Mechanism	Alcubierre Bubble	Comparative Advantage
Principle	Matter phase transition (changes the actor)	Spacetime deformation (changes the stage)	Revoredo: philosophically deeper
Domain	Hyperspace (multidimensional bulk)	Conventional spacetime	Revoredo for very long journeys
Required energy	Planck scale via ZPE	Negative mass equivalent to Jupiter	Both require Kardashev II+
Proposed source	Dyson Sphere + ZPE (ER=EPR)	Undefined; exotic matter not observed	Revoredo: more concrete source
Optimal Kardashev scale	Type II–III (intergalactic)	Type I–II (local interstellar)	Complementary, not rivals
Causality	Orthogonal time; paradoxes mitigated	Causal violations unresolved	Revoredo: more sophisticated handling

Beyond the Veil: Uncovering the Truth About UFOs

Crew during travel	Practically instantaneous subjectively	Normal time inside bubble	Different depending on operational need
--------------------	--	---------------------------	---

A2.14.2. The Revoredo Mechanism as an Intergalactic Backbone Network

Since the tachyonic state theoretically travels at infinite speeds with zero energy once conversion is complete, the Revoredo Effect would be the equivalent of high-speed intergalactic fiber optics: it would connect galaxy clusters and principal stellar systems. The Dyson Spheres at points of origin would act as quantum energy routers, maintaining entanglement with ships in hyperspace and simultaneously serving as beacons of the Hyperspace Positioning System, while maintaining the entangled counterpart of the $\text{PMT}^{\text{target}}$ for each ship in transit.

A2.14.3. The Alcubierre Metric for Local Navigation and Exploration

Once the ship uses the Revoredo Mechanism to arrive instantaneously at a sector and reverts its phase to normal matter, it would employ the Alcubierre Engine to maneuver within that stellar system. Since Alcubierre deforms real space, it is ideal for tactical maneuvers, planetary mapping, and approaches where jumping into tachyonic hyperspace would be excessive or dangerous due to the absence of a stable local entanglement station.

A2.14.4. The Technological Signature in the Cosmos

A civilization using both engines would leave two potentially detectable cosmic signatures:

- **Alcubierre signature:** massive gravitational waves and high-energy Hawking radiation bursts when the bubble halts.
- **Revoredo signature:** extreme thermal anomalies in stars (ZPE extraction by the Dyson Sphere) combined with neutrino bursts or remnant particles from tachyon

Beyond the Veil: Uncovering the Truth About UFOs

condensation at the precise coordinates where ships re-enter the three-dimensional brane.

A2.15. Conclusions and Epistemological Status of the Model

The Revoredo Model constitutes an internally consistent body of theoretical physical hypotheses, resting on the same mathematical pillars that underpin string theory, the Alcubierre metric, and the ER=EPR conjecture: algebraic consistency, logical extension of existing formal frameworks, and absence of internal contradictions.

A2.15.1. Epistemological Status of the Revoredo Model

Status: speculative theoretical physics hypothesis with internal mathematical consistency.

Mathematical basis: General Relativity (tensors), Quantum Field Theory (KG with $m^2 < 0$), String Theory (vibration modes), Differential Geometry (Calabi-Yau).

Current verifiability: not experimentally verifiable with present technology (requires Planck scale).

Epistemological comparison: identical status to the Alcubierre metric, the ER=EPR conjecture, bosonic string theory, and cosmic inflation models. This status refers to the type of hypothesis: a speculative theory with internal mathematical consistency.

Falsifiability: the model is in principle falsifiable under a unified theory of quantum gravity that determines the behavior of spacetime at the Planck scale.

Own postulates: Within the model, most of the pieces come from established physics or from respected, actively researched theories. The model's own postulates, neither proven nor ruled out, are: the extraction of zero-point energy (A2.7), nonlocal energy compensation with the Dyson Sphere, the closing of strings by the torsion field (A2.6.2), the inhibited condensation sustained by the engine (RIC), the restoration of the PMT with the inverted Lindblad term, the action of time reversal on μ in hyperspace (A2.6.4.2), and the encoding of the

Beyond the Veil: Uncovering the Truth About UFOs

bulk's geometry in the entanglement network (A2.12.3). They presuppose a Type II-III civilization and engineering that, if possible, lies centuries away.

The central conceptual leap of the Revoredo Mechanism over the Alcubierre metric is philosophically profound: while Alcubierre proposes modifying the stage (spacetime) to transport an actor without changing its nature, Revoredo proposes modifying the actor (matter) so that the stage becomes irrelevant. This distinction is not merely physical but ontological.

The derivation of the Phase Memory Tensor from first principles (chain $\varrho \rightarrow W(x,p) \rightarrow C_{ab}(x) \rightarrow M_{\mu\nu}$, dynamic evolution equation, conservation law, and fidelity condition) closes the most important formal gap in the model: how to preserve the identity of an object across a quantum phase transition at the Planck scale. This derivation rests on real mathematical objects used in their correct speculative extension, with the same level of rigor as any other hypothesis in advanced theoretical physics.

The history of physics teaches humility before barriers declared impassable: the sound barrier was an aeronautical limit, not a natural one. The quantum foam of spacetime at the Planck scale is today terra incognita. In that unknown territory, the Revoredo Model plants a mathematically coherent and structurally complete hypothesis.

A2.16. Conceptual References

The following theories, equations, and formal frameworks constitute the foundations on which the Revoredo Model rests:

- Alcubierre, M. (1994) — Alcubierre metric: spacetime deformation for apparent superluminal propulsion. Direct reference of comparison.
- Bender, C.M. & Boettcher, S. (1998) — “Real Spectra in Non-Hermitian Hamiltonians Having PT Symmetry,” Phys. Rev. Lett. 80, 5243. Foundation of PT-symmetric quantum mechanics used in A2.6.4.
- Bender, C.M., Brody, D.C. & Jones, H.F. (2002) — “Complex Extension of Quantum Mechanics,” Phys. Rev. Lett. 89, 270401. Construction of the CPT inner product and unitary evolution under non-Hermitian PT-

Beyond the Veil: Uncovering the Truth About UFOs

symmetric Hamiltonians, used for the Revoredo Metric Operator.

- Bender, C.M., Brody, D.C. & Jones, H.F. (2003) — “Must a Hamiltonian be Hermitian?” *Am. J. Phys.* 71, 1095. General review of PT-symmetric quantum mechanics and its physical interpretation.
- Calabi, E. & Yau, S.T. — Calabi-Yau spaces: geometry of compactified dimensions in string theory.
- Casimir, H.B.G. — Casimir effect: quantum pressure between conducting surfaces; basis of the ZPE extraction mechanism.
- Dyson, F. — Dyson Sphere: stellar energy-capture megastructure; reinterpreted as quantum infrastructure, energy router, and PMT^{target} repository.
- Einstein, A. — Special and General Relativity: relativistic energy equation, energy-momentum tensor, Christoffel symbols, geodesics, vierbein.
- Green, M., Schwarz, J. & Witten, E. — String Theory: vibration modes as the origin of particles, open and closed strings, bosonic tachyonic strings.
- Heisenberg, W. — Uncertainty Principle: foundation of quantum vacuum fluctuations.
- Higgs, P. et al. — Spontaneous symmetry breaking mechanism (Mexican hat): mathematical analog of tachyon condensation.
- Kaluza, T. & Klein, O. — Theory of compactified extra dimensions: foundation for the existence of hyperspace.
- Kardashev, N. — Kardashev scale: classification of civilizations by energy consumption; framework for the model's applicability.
- Klein, O. & Gordon, W. — Klein-Gordon equation: basis of the tachyonic formulation in QFT with $m^2 < 0$.
- Lindblad, G. — Lindblad master equation: open-system equation formalism; basis of the PMT dynamic restoration term (operating in reverse, as anticipation).

Beyond the Veil: Uncovering the Truth About UFOs

- Maldacena, J. & Susskind, L. — ER=EPR conjecture: equivalence between quantum entanglement and micro-wormholes. Basis of the nonlocal energy compensation channel, the Hyperspace Positioning System, and the $J_{\mu\nu}^{\text{EPR}}$ term of the dynamic PMT.
- Sen, A. — Tachyonic matter and string condensation: equation-of-state formalism for the tachyonic field in the superluminal cruise regime.
- Sen, A. (2002a) — “Rolling Tachyon,” JHEP 0204:048. Derivation of the tachyon effective action and rolling solution used in A2.4.3.
- Sen, A. (2002b) — “Tachyon Matter,” JHEP 0207:065. Asymptotic pressureless “tachyon matter” regime used for the Tachyonic Saturation stage.
- von Neumann, J. — Density matrix evolution equation ($\partial_t \varrho = -i[H, \varrho]/\hbar$): basis of the unitary evolution term of the dynamic PMT.
- Wigner, E. & Weyl, H. — Wigner function and Weyl transform: mathematical bridge between Hilbert space and phase space; basis of the PMT projection into spacetime.
- Wilczek, F. — Time Crystals: structures with spontaneous temporal periodicity; proposed as phase stabilizers and physical repositories of $\text{PMT}^{\text{target}}$.